

COLUMBIA LIBRARIES OFFSITE
HEALTH SCIENCES STANDARD



HX64057666

RD33 B76 1907

Five hundred surgica

RECAP

500

SURGICAL SUGGESTIONS

WALTER M. BRICKNER, M.D.

ELI MOSCHCOWITZ, M.D.



COLLEGE OF PHYSICIANS AND SURGEONS
COLUMBIA UNIVERSITY



1545 GASPAR TALIAACOTIUS 1599

THE JEROME P. WEBSTER
LIBRARY OF PLASTIC SURGERY

ALUMNA UNIVERSITY



500

SURGICAL SUGGESTIONS

500

SURGICAL SUGGESTIONS

PRACTICAL BREVITIES IN DIAGNOSIS AND TREATMENT

BY

WALTER M. BRICKNER, B. S., M. D.

*Chief of Surgical Department, Mount Sinai Hospital Dispensary;
Editor-in-Chief, American Journal of Surgery,
New York,*

AND

ELI MOSCHCOWITZ, A. B., M. D.

*Assistant Physician, Mount Sinai Hospital Dispensary;
Associate Editor, American Journal of Surgery,
New York.*

SECOND SERIES



NEW YORK, U. S. A.
SURGERY PUBLISHING COMPANY
92 WILLIAM STREET
1907

~~W. H. Lister Library~~

RD 33

B76

1907

Copyright, 1907
BY
SURGERY PUBLISHING COMPANY

PREFACE

The first issue of "Surgical Suggestions," published in 1906, was exhausted in a few months. The continued demand for the little book has led us to prepare this second series. In it we have incorporated with the "Suggestions" that were in the first edition, as many more, making the total number about five hundred.

These practical brevities make no pretensions to completeness. They are merely observations taken here and there from our own surgical experiences.

W. M. B.
E. M.

August, 1907

SURGICAL SUGGESTIONS.



In determining whether or not to operate after injuries to the head, a surgical judgment of the case is usually better than one based strictly on the application of neurological rules. Cranium.

There is no class of cases in which a prognosis is so often at variance with the extent of the injury as in cranial injuries. The prognosis in such cases should, therefore, always be guarded.

In cases of head traumata, bleeding from the mouth or nose does not necessarily mean that the case is one of fracture at the base. The hemorrhage may be entirely due to a localized injury.

In fractures of the base of the skull with bleeding from the ear it is necessary to keep the auditory canal absolutely clean in order to prevent infection of the meninges.

“Egg shell crackle” elicited in palpating a tumor of the cranial bones is diagnostic of sarcoma originating in the diploe.

HEAD.

Depilatories are useful in the preparation of the scalp for the treatment of abscesses or infected wounds, when the nature of the infection or the matted condition of the hair makes shaving difficult.

Brain. Tumors of the brain frequently simulate, in their earlier stages, diseases of the stomach.

Lumbar puncture must not be performed in cases of tumor of the brain. Sudden death has frequently happened in such cases.

In exploring for tumors of the brain, the best guide for determining an isolated hardness is the finger; the use of a needle is very deceptive.

Ear. A furuncle deeply situated in the external auditory canal gives signs that may be mistaken for mastoiditis. Great pain when the concha is moved about, will serve to differentiate it from the latter.

Severe and repeated headaches may be due to the unsuspected presence of otitis media, with or without mastoiditis.

In cases of unaccountable fever, especially in children, never fail to examine the ear.

Tinnitus aurium, present only in the recumbent posture, is suggestive of aneurism of one of the posterior cerebral vessels.

The history of a discharge from an ear appearing a few days to a few weeks after the beginning of a slowly developing deafness in that ear, unaccompanied at any time by pain, is suspicious of tuberculous otitis media.

A bean-shaped pulsating swelling just below the mastoid apex, in cases of mastoiditis, may be only a lymphatic gland, but it may also be a thrombosed jugular vein. Its nature should therefore be determined before the operation is concluded.

During mastoid operations always sever with scissors any fragment of tissue attached to a bit of bone loosened with the chisel or rongeur, before removing it. The tearing out of a fiber of the sterno-mastoid muscle, for example, will open a channel of infection in the neck.

HEAD.

If the zygomatic cells are thoroughly laid open, one frequent cause of persistent suppuration requiring secondary mastoid operation, may be avoided.

A persistent elevation of temperature after a radical operation for mastoiditis should lead one to suspect the possibility of a complicating brain abscess. If the fever shows wide fluctuations of temperature a sinus thrombosis is more probably the cause.

Streptococcic infections having their entrance through the ear are more apt to affect the muscles than similar infections with other points of entry.

Nitrate of silver may be attached in full strength to the end of a probe, as for application in the middle ear, by heating the tip of the instrument and pressing it into the stick of caustic; a little of the latter will melt and form a bead on the probe when it cools.

Eyc. Cystic swellings at the external angle of the eye are usually dermoids. In some cases they communicate by a small opening with an intracranial sac.

HEAD.

After the extraction of a foreign body from the cornea, a drop of castor oil between the lids will ameliorate the pain.

Severe pain in the orbit or even in the eye itself should make one think of frontal sinus infection, especially if there is, or recently has been, a nasal discharge. Marked localized tenderness will soon confirm the suspicion, if the disease exist.

Nose.

A diagnosis of supraorbital neuralgia should not be made until frontal sinusitis has been carefully excluded.

A persistent, chronic discharge from the nose should lead one to suspect chronic disease of the frontal or other accessory sinus.

Transillumination is a method of corroborative value only in the diagnosis of accessory nasal sinus disease. By itself it is of small diagnostic use.

Frontal sinus suppuration rarely requires a disfiguring operation for its relief. It can usually be satisfactorily dealt with through an opening in the line of the eyebrow.

HEAD.

In every case of injury to the nose, with or without fracture, it is well to examine the septum for displacement. If displaced it should be carefully restored, using a nasal plug, if necessary, to keep it in place.

Face.

Non-malignant tumors of the parotid practically never cause pressure effects on the facial nerve. This may be of importance in differentiating them from malignant tumors.

A swelling in the parotid region is not necessarily a part of the parotid gland. It may be an infection of the pre-auricular lymphatic gland. Such an enlargement may be associated with herpes of the forehead, or, sometimes, it may be part of a chain of tuberculous lymph glands.

The position of Steno's duct must be remembered when operating on the face.

In chronic osteomyelitis of the jaw it is better to wait months for a sequestrum to form than to operate a dozen times for the removal of necrosed bone.

All swellings of the lower jaw accompanied by discharging fistula, especially mul-

HEAD.

tiple fistulæ, should be looked upon with the suspicion of actinomycosis until proven to be otherwise.

If a frightened or refractory child will not open its mouth, pass a probe between two teeth and back to the palate. Instantly the mouth will open and a gag may be slipped in.

Mouth and Pharynx.

The employment of adrenalin as an application with cocain to the mucous membrane of the cheek, e. g., for the excision of a leukoplakic ulcer, is not to be advised. There may be severe secondary hemorrhage.

When there is bleeding from the tongue, post-operative or otherwise, and one feels reasonably sure that the hemorrhage is arterial, it can, as a rule, be easily arrested by passing the forefinger down to the epiglottis and hyoid bone and drawing the base of the tongue upward toward the chin.

It is wrong to perform any radical operation for an ulcer of the tongue without preliminary microscopical examination. Clinical symptoms, no matter how typical, are often misleading.

HEAD.

A deep ulceration of the fauces or tonsils should not be diagnosed as specific without excluding acute lymphatic leukemia.

Before operating for pharyngeal adenoids or hypertrophied tonsils make sure that these are not merely an expression of status lymphaticus. If they are, do not employ an anesthetic. Also determine whether the patient is a hemophiliac. If he is, do not operate at all.

When opening a retropharyngeal or peritonsillar abscess in a small child, by the buccal route, have the head dependent and instruments at hand for tracheotomy. These instruments are needed but rarely, but then urgently.

NECK.

In seeking a cause for torticollis, don't fail to examine the teeth.

In all cases of torticollis, examine for caries of the spine.

Pediculosis capitis may be the indirect cause of acute torticollis by reason of a developing post-cervical adenitis.

A submaxillary swelling should not be dismissed as a lymphatic adenitis without studying Wharton's duct on the same side. Massage of pus therefrom would alter that diagnosis.

Examination into the nature and cause of discrete hard lymphatic swellings on each side of the neck, along the sterno-mastoid, should include exploration of the pharynx and naso-pharynx for possible new growth.

In all glandular affections of the neck it is quite as important to treat the source of infection, e. g., carious teeth, as to treat the inflamed glands.

In "Ludwig's angina," the cardinal principle in the treatment is extensive incision. An incision that passes no matter how deep into the substance of the submaxillary gland proper, will prove of little avail unless the tissues within the wound have been broken up until they are practically pulpy.

Have the tracheotomy instruments handy before operating upon a case of angina Ludovici.

NECK.

The greatest ultimate danger in cut-throat cases is the onset of a septic pneumonia. This may be obviated in a measure by closing up the pharyngeal wall, and by paying the strictest attention to asepsis.

In cut-throat wounds where the thyro-hyoid membrane has been severed, it is necessary, in order to restore perfect phonation and deglutition, to suture this membrane accurately.

Avoid the use of peroxid of hydrogen in wounds of the neck. It is too apt to dissect up the loose cellular planes. The same warning applies in many cases of cellulitis of the hand or foot.

In the presence of a hard, diffuse, chronic swelling in the neck having some of the appearances of a malignant growth, the possibility that the tumor is a so-called "woody phlegmon of Reclus" must be considered.

Hard subcutaneous tumors of the upper third of the neck, with signs of malignancy, are often epitheliomata arising from branchial clefts.

In performing operations on the neck, make the skin incision parallel to the muscular plane.

Do not empty a thyro-glossal cyst by aspiration before extirpating it. It is well to inject the cavity with a methylene blue solution first, in order to make sure that all parts of the cyst wall will be extirpated. Another method is to first empty the cyst and then fill it with paraffin.

In all operations in the left subclavian triangle of the neck, the location there of the thoracic duct must not be forgotten.

The best thing to do in such emergencies as air embolism is to apply compression immediately and pour large quantities of solution, preferably salt solution, into the wound.

Gradually increasing hoarseness in people past middle age, without definite cause, and with a history of pain radiating to the ear, is suggestive of malignancy.

NECK.

Tracheotomy. In urgent cases a high tracheotomy should be performed, not a low tracheotomy. The former can be done very rapidly; the latter requires considerable dissection.

The best site for an urgent tracheotomy is through the crico-thyroid membrane. To hold the opening apart a couple of hairpins, bent at the end, may be used as retractors.

In the performance of high tracheotomy a great deal of room can be gained by dividing transversely the fascia that extends upward from the thyroid.

After tracheotomy the air of the patient's room should be kept reasonably warm and moist. Draughts of cold air provoke much irritation.

The greatest immediate danger after a tracheotomy is the possibility of a subsequent pneumonia. This can, in a large measure, be obviated by filtering the inspired air through a soft sponge saturated with warm one per cent. phenol solution.

NECK.

Repeated attacks of coughing after tracheotomy may mean irritation of the posterior wall of the trachea by the tube; change the length or shape of the canula.

BREAST.

A "tumor" of the breast occasionally proves to be only a chronic abscess. It has happened that a breast amputated for carcinoma has been found to be the seat of old abscesses only.

In the treatment of a breast abscess the size of the incision is not as important as its location and direction.

A small incision and the proper employment of Bier's breast cup will secure exceedingly gratifying results in the management of breast abscesses.

In the presence of a breast infection that fails to heal within a reasonable time after appropriate incision and dressings, it is well to think of local tuberculosis.

The appearance of pus in the breast of a woman who is not, or has not recently been nursing, is suspicious of some unusual form of infection, e. g., tuberculosis.

BREAST.

A tender, painful swelling just at or beyond the upper, outer border of the breast, and near the edge of the pectoralis major, is usually an inflamed lymphatic gland. In its presence it is well to look for some skin infection about the waist line, e. g., furuncles, which are not rare at this site as a result of irritation by the corset. Per contra, with a boil, abscess, dermatitis or other infection at or above the waist line, one may be on the lookout for glandular enlargement at the point referred to.

In the performance of the radical operation for breast carcinoma it is important to avoid injury to the periosteum of the ribs.

Multiplicity of tumors of the breast usually speaks against carcinoma.

Breast tumors, especially in the early stages, not seldom fail to present the classical signs by which their malignancy or non-malignancy may be determined clinically. In all cases where the diagnosis is doubtful, a specimen of the tumor should be removed for microscopic study, before undertaking a radical operation.

In strapping the chest for fractured rib, two points should be particularly noted: 1. The straps should pass well beyond the median line. 2. They should be applied in full expiration. One or two straps passed over the shoulder help much to secure immobilization.

Cold abscess and lipoma often simulate each other very closely, especially around the chest. If in doubt, aspirate.

Do not be too hasty in making a diagnosis of intercostal neuralgia. With the exception of pulmonary and pleural conditions, ulcer of the stomach simulates intercostal neuralgia more frequently than any other lesion.

It is remarkable how frequently a purulent pericarditis may exist without causing many or severe symptoms. Never neglect an examination of the cardiac area, therefore, in cases of suspected sepsis.

When a patient complains of dysphagia, do not neglect to examine the pericardium for effusion.

THORAX.

A history of attacks with symptoms of esophageal stricture and intervening periods of well-being is suggestive of cardiospasm.

A satisfactory method of x-ray study of esophageal diverticula and strictures consists in fluoroscopy of the thorax while the patient is swallowing an emulsion of bismuth subnitrate. A skiagraph may be made immediately afterwards as a supplementary record.

Before operating for sarcoma examine the lungs carefully. Do not operate if the patient has persistent cough and blood-stained sputum (not due to tuberculosis), even though no definite signs are found in the lungs—a metastasis has developed.

A fluctuating swelling appearing between the ribs may, of course, be tuberculous or syphilitic in origin, but it may also be an extension of an intrathoracic growth, e. g., dermoid cyst of the mediastinum. In all such cases, therefore, a careful examination, by auscultation and percussion, should be made.

A slender fish bone lodged in a bronchus will usually not cast a shadow on the x-ray plate. In such a case bronchoscopy and auscultation are more reliable diagnostic measures. In addition to a variety of moist rales, one may hear, associated with the inspiratory or expiratory murmur, or both, a musical or vibratory note, when a bone or pin lies in a bronchus.

If a patient dates irregular or persistent cough from a time when he thinks he "swallowed" or inspired a foreign body, the fact that the physical signs elicited upon examination of the chest are peculiar—different from those found in ordinary types of bronchitis—points strongly to the presence of a foreign body.

Bronchiectasis is not seldom complicated by brain abscess.

In aspirating the chest, see to it that the syringe is in good condition before inserting the needle. Never apply the syringe to the needle after the latter has been inserted; a severe pneumothorax may result. If the syringe is found to be out of order while the aspiration is being done, withdraw the needle also and reinsert.

THORAX.

There is one point that must always be thought of when pus has been aspirated after an exploratory puncture for either suspected empyema or liver abscess,—to make sure that the “pus” does not come from a bronchus. This can be determined, as a rule, by microscopical examination of the aspirated fluid.

Very extensive and rapidly spreading subcutaneous infections may result after an aspiration of a foul-smelling empyema. It is therefore wise to always operate over the site of aspiration, and especially to see that the puncture wound is well drained.

When operating for empyema thoracis it is a good rule to aspirate again when the pleura is exposed and before it is incised. This may save some embarrassment.

The shock of evacuating an empyema thoracis may be largely avoided by making but a small opening in the pleura (after resecting the rib) and applying at once several thicknesses of gauze. At the next dressing much or most of the pus will be found to have escaped into the gauze, and the pleural wound may then be enlarged without producing shock.

THORAX.

After operation for empyema, a cover of oiled silk or gutta-percha over the gauze dressing serves to prevent admission of air into the pleural cavity, while it will not interfere with the escape of air already in the chest. Indeed, a flap of rubber may be laid over the wound and fastened with a little chloroform above. This allows pus to escape from beneath it and excludes the admission of air.

ABDOMEN.

It is surprising how much information can be derived by abdominal palpation conducted with the patient in a hot bath, the temperature of the water being gradually raised to 105° F. It usually secures as much relaxation as does the administration of an anesthetic, sometimes even more. In addition to the avoidance of the dangers and the disagreeable features of narcosis, it has the important advantage that the patient is able to call the examiner's attention to sensitive areas.

In the presence of a tumor in the mid-line between umbilicus and pubes, the possibility of a cyst of the urachus must be borne in mind. It may simulate an ovarian cyst or other tumor, or a distended bladder.

ABDOMEN.

Eczema of the umbilicus is sometimes merely the expression of an infected dermoid cyst at that site.

A discharge from the umbilicus may be due to an infected dermoid cyst, to an eczema of the umbilicus, to a patent urachus (urine), to a cyst of the urachus (milky discharge); it may be of pus from an abscess within the abdomen or in the abdominal wall, or of feces (Meckel's diverticulum, perforated strangulated hernia, fecal abscess from tuberculosis).

Do not ligate tumors of the navel without making sure that the intestine is not included within the ligature.

In performing paracentesis in the median line for abdominal fluid, be sure that the bladder is empty. When it is necessary to perform paracentesis in the lateral part of the abdomen, be careful to avoid the deep epigastric artery.

Unless some other cause is evident don't fail to examine for signs of tabes when an adult complains of pains about the waist, in the back or in the lower extremities.

Children who complain frequently of pain in the stomach should be examined for evidence of beginning Pott's disease. Such cases, treated before the development of curvature, usually yield very satisfactory results.

In all cases of acute abdominal pain, never fail to examine the lungs and gums. The onset of pneumonia or pleurisy frequently closely simulates acute appendicitis; lead colic may simulate almost any painful abdominal condition.

Enlargement of the veins at the side of the abdomen is indicative of obstruction to the flow of blood in the inferior vena cava; distention of veins about the umbilicus suggests obstruction in the portal circulation. The former may be associated with varices of the lower extremities, the latter with hemorrhoids.

In all cases of recurrent vomiting examine the midline of the abdomen for a small epigastric hernia.

Catheterization sometimes makes the evidences of "appendicitis" or "abdominal tu-

ABDOMEN.

mor" vanish with the escape of the urine from a distended bladder.

In cases of run-over by vehicles, if the wheels pass over the trunk from right to left the liver is the organ most commonly ruptured, whereas, if the wheels pass from left to right the spleen is more frequently injured.

The hypodermatic injection of eserine (salicylate) gr. $1/30$ — $1/40$, during or just after an abdominal operation, will, in most cases, entirely or largely prevent the distressing tympanites that otherwise usually occurs.

A pulsating swelling in the midline of the abdomen should not be too quickly accepted as an aneurism of the aorta. It may be a retroperitoneal tumor.

In the presence of large masses of glands in the epigastrium, especially on the right side, examine the testicles for new growth.

Large, slowly growing, slightly movable abdominal tumors near the median line, caus-

ing few symptoms and not accompanied by signs of malignancy, are suggestive of mesenteric cysts.

A primary tumor of the lateral abdominal region in infants and young children is usually a sarcoma of the kidney.

In operating for perforating gun-shot wounds of the abdomen, find the source of any bleeding first, before attempting to suture any perforation.

In suturing the fascial layers of the abdominal wall do not take too large bites with the needle. Necrosis may occur, and sloughing of the fascia predisposes to the formation of hernia.

If there is repeated vomiting and the patient shows some evidences of collapse, after a laparotomy, especially after operations in the gastric region, examine for separation of the wound and prolapse of the abdominal contents.

In children, in cases of peritonitis of unknown origin, examine for gonorrheal vulvovaginitis.

ABDOMEN.

Steady loss of weight without other demonstrable cause should lead the physician to look for a possible malignant visceral neoplasm. Persistent "indigestion" due to some condition not positively ascertained, should be submitted to surgical diagnosis.

Bile Tract. Repeated attacks of "indigestion," not obviously due to some other condition, should awaken the suspicion of gall-stones. Most of the patients operated upon for cholelithiasis give a history of having been treated for a long time for "dyspepsia," and in many of these cases the correct diagnosis might earlier have been established.

If pressure in the right hypogastrium gives rise to a referred pain in the shoulder region, the offending area is probably the gall-bladder and not the pylorus.

In an attack of cholelithiasis the vomiting as a rule is not attended by relief of pain; the contrary is true in ulcer of the stomach.

In differentiating between gastric ulcer and gall-stone pains, the association of a chill usually points to cholelithiasis.

Tuberculosis and cholelithiasis are only very rarely associated.

Long pauses between attacks of gastric or abdominal pain speak in favor of cholelithiasis.

In the progress of a cholecystectomy, if a stone slips away after cutting through the cystic duct, and cannot be found, no great anxiety need be felt, for the stone usually comes away spontaneously in the subsequent discharge.

Great pain following any operation upon the biliary tract should always lead one to suspect leakage of bile into Morrison's space. If such should be found to be the case insert a drainage tube.

When operating for cholelithiasis, don't fail to examine the hepatic duct.

When palpating the common bile duct for stone, make sure that a suspected calculus is not a gland.

In catarrhal icterus the pulse is usually slow; in jaundice from cholelithiasis this is usually not the case.

ABDOMEN.

Gradually increasing jaundice without previous history of pain, or with a history of very slight pain, is very suggestive of malignant disease.

Before attributing enlargement of the liver to a surgical condition exclude chronic hepatic congestion of cardiac disease.

Examine the rectum in all cases of tumor of the liver. Likewise, before operating for cancer of the rectum examine the liver for metastasis.

Stomach.

Examination to determine the possible presence of cardiac disease or aneurism should always be made before passing a stomach tube. In the presence of such lesions the tube should not be employed except as a life-saving measure in an emergency.

A hasty diagnosis of ulcer of the stomach should not be made merely because the patient has vomited suddenly large quantities of blood. If the bleeding occurs at regular intervals the possibility of vicarious menstruation must be considered.

It is well to remember that not all ulcers of the stomach are characterized by the classical symptoms of pain, vomiting and hemorrhage. Many patients presenting "dyspeptic" symptoms of only mild grade are afflicted with this disease, and such cases may easily be diagnosed as functional disorders until the persistence of the symptoms leads one to suspect the graver malady.

The thirst following a hemorrhage from gastric ulcer is best relieved by small quantities of cocain in solution.

If an undoubted case of ulcer of the stomach is associated with chills, in most cases it means that the ulcer is adherent to the spleen.

A sudden desire for sharp, sour and spicy articles of food in a middle-aged or elderly person is often the first symptom of a beginning gastric carcinoma.

In a case of gastric disease of doubtful diagnosis, progressive loss of weight is the most important sign in determining the probability of carcinoma.

ABDOMEN.

The possibility of gastric cancer must be considered in cases of supposed pernicious anemia.

In persons of middle age presenting gastric symptoms, the diagnosis of cancer should not be excluded because the symptoms have had a sudden onset. Such an onset occurs in a fair proportion of cases.

Vomiting, secondary anemia and absence of free hydrochloric acid in the gastric juice—a triad of symptoms at once suggestive of carcinoma ventriculi—may occur as a result of chronic nephritis. Especially if the urine contain no casts or albumin is the observer apt to be led astray.

If a patient vomits coffee-ground material in which no lactic acid is present, one can almost always exclude carcinoma.

A reasonable suspicion of the presence of a cancer in the stomach or intestine is sufficient indication for explorative operation.

Before proceeding with a radical operation for carcinoma of the stomach, examine not

ABDOMEN.

only the liver but also the general abdominal cavity, especially the pelvis, and in females the ovaries, for any sign of metastasis.

Do not be too sure that a mass in the region of the pylorus is a carcinoma. In some cases the infiltration around a chronic ulcer is very extensive and may simulate the feel of a new growth.

If a patient begins to vomit long after a radical operation for carcinoma of the stomach, do not jump to the conclusion that the cause is a local recurrence. It may be a metastasis in the brain.

In a patient with spondylitis, symptoms simulating acute peritonitis may be due to acute dilatation of the stomach.

The occurrence after laparotomy of marked distention of the upper abdominal zone, vomiting and collapse points to acute dilatation of the stomach.

It is a peculiar fact that post-operative prolapse through the epigastric wound occurs frequently in operations for malignant disease

ABDOMEN.

of the stomach. Such wound therefore should be closed with more than usual firmness and all possible precautions should be taken to guard against post-operative vomiting.

In performing posterior gastro-enterostomy see that the opening in the transverse mesocolon is not so narrow that it may constrict the anastomosed segment of small intestine nor yet so large that it may permit of a possible hernia into the lesser sac. By inserting a number of sutures between the mesocolon and the stomach wall about the anastomosis these possibilities may, in large part, be obviated.

If within a week or two after the performance of gastrostomy the drainage tube should be expelled from the fistula, do not entrust its re-introduction to inexperienced hands. It has sometimes happened that the tube has been pushed into the peritoneal cavity, instead of into the stomach.

Intestines. In the presence of anemia or of faintness, without other apparent cause, inquire concerning the passage of black stools. The condition may result from hemorrhages due

to an ulcer or neoplasm of the small intestine.

A gradually increasing anemia in an elderly person, without any other symptoms, is highly suggestive of a latent carcinoma, often in the intestine.

In typhoid fever spontaneous rupture of the spleen may simulate intestinal perforation.

The triad of symptoms—pain, vomiting and distention—without fever, points to intestinal obstruction.

Attacks of abdominal pain preceded by “rumbling” of the bowels is suggestive of some obstructive condition.

The passage of gas or even of a small amount of feces, after an enema, does not gainsay the presence of intestinal obstruction.

In an acute condition simulating intestinal obstruction, if a large mass can be felt in the abdomen think of omental torsion.

ABDOMEN.

An attack of acute intestinal obstruction, with passage of blood, and in the presence of a cardiac lesion, is suggestive of thrombosis of a mesenteric vessel.

Every case of intestinal obstruction of obscure origin should be inquired into closely with reference to a previous history of cholelithiasis. If a definite history of this is obtained, it is well to suspect obstruction by a gall-stone.

When operating for volvulus of the large intestine, insert a rectal tube as high up as possible before attempting the reduction. The volvulus will quickly collapse and the necessity for evisceration will thus be avoided.

In infants sudden, severe colic associated with diarrhea or the passage of small quantities of blood, should lead one strongly to suspect an intussusception.

When reducing an intussusception don't pull on the intussusceptum but push on the intussusciens.

Whenever we meet with a movable, sausage-shaped mass in the abdomen, with a history of chronicity, it is well to think of the possibility of its being a case of hyperplastic tuberculosis of the intestine. This diagnosis will be rendered more suggestive if there are definite signs in the lungs.

Simple or multiple enterostomy, usually with prompt suture of the opening, is many times a life-saving operation in the presence of intestinal paresis, as from general peritonitis.

At the onset of an attack of acute appendicitis the pain is usually referred to the gastric region.

Appendix.

The cessation of severe pain during the course of acute appendicitis often means perforation.

The twisting of the pedicle of a small ovarian cyst may simulate both the symptoms and signs of attacks of appendicitis.

The tenderness in appendicitis may not be (probably usually is not) just at McBur-

ABDOMEN.

ney's point. The base of the appendix is, however, usually at, or near, that point. The site of greatest tenderness is often over the tip of the appendix. A line drawn between that site and McBurney's point will many times represent the general direction in which the appendix is lying.

In cases of chronic appendicitis, if an examination be conducted with the patient in a hot bath (105° F.), the thickened appendix may often be felt to roll under the finger.

In a tuberculous patient with supposed chronic appendicitis it is well to suspect tuberculous disease of the ileo-cecal valve.

Hernia. It is a wise rule never to attempt taxis in cases of strangulated hernia. The only exception might possibly arise in a case seen within the first hour.

All cases of hernia in which there is a history of frequent urination should lead one to the suspicion that the hernial sac contains part of the bladder.

ABDOMEN.

If a peculiar looking mass is found at the inner side of the ring in the course of an operation for inguinal hernia, do not incise or dissect it before convincing yourself that it is not the bladder.

Do not be too hasty in resecting a strangulated loop of intestine. It is remarkable how frequently such loops become viable after long continued applications of hot saline solution.

Examine the umbilicus and the inguinal and femoral canals in all cases of obscure intestinal obstruction. Small strangulated femoral herniæ often simulate very closely the feel and appearance of a gland, and in such cases one may be easily misled.

In ligating the omentum, it is a good rule never to place a ligature around a piece larger than the width of a finger.

Hemorrhage from the bowel in children is not infrequently caused by a polypus in the rectum.

RECTUM.

Prolapsus ani is a frequent accompaniment of bladder stone in children.

RECTUM.

Prolapse of the rectum in children usually yields to treatment by strapping the nates together with adhesive plaster, if carried out intelligently and persistently, for several weeks or months. The child should be obliged to defecate in the recumbent posture and while the strap is on. After defecation the strap is removed, the parts cleansed and a fresh strap applied, all while the child is recumbent.

A mass protruding from the rectum of an infant or child may be an intussusception and not a mere prolapse.

Don't fail to make a digital rectal examination in cases of appendicitis and in all ailments when the diagnosis is obscure. Nor should it ever be omitted before an operation upon anal disorders. It may save the embarrassment of a subsequent discovery that a patient's hemorrhoids, for example, were but an expression of a carcinoma higher up in the rectum.

A radical operation for hemorrhoids should not be undertaken until the etiology of the piles has been determined. Some-

times the cause is an obstruction in the portal circulation due to hepatic disease. Per contra, abscess of the liver may be due to infection from a hemorrhoid operation performed even some months before.

Bleeding from capillary hemorrhoids high in the rectum usually yields to injections of cold water, or a cold solution of tannic acid. In these cases, however, it is important to exclude the presence of an ulcer further up.

If a thrombosing pile is opened before the clotting is complete, it is very apt to fill up again and may even become edematous and inflamed.

When removing hemorrhoids much after-pain may be obviated by making radiating nicks in the skin margin of the anus.

After an operation for hemorrhoids it is desirable to insert into the rectum a tampon canula, made by smearing with vaselin gauze layers wrapped about a piece of rubber tubing, about three inches long and trans-fixed at its distal extremity with a large safety pin. The tampon canula prevents

RECTUM.

oozing by its gentle pressure, allows any considerable hemorrhage to show itself externally, makes the escape of flatus painless and the introduction of an oil enema easy.

Although profound anesthesia is required to abolish the anal reflex, chloroform or ether is not always needed in order to divulse the sphincter ani. This may be accomplished painlessly, and usually with fair satisfaction, under ethyl chlorid or nitrous oxid narcosis if, especially, an opium suppository is introduced a half-hour beforehand, and a pledget of cotton wet in cocain solution is applied just before the operation.

After operations upon the rectum, especially after those involving divulsion of the sphincter ani, voluntary urination is apt to be inhibited for a day or more. This is especially the case when stretching is done in a sagittal direction, i. e., towards the urethra and the coccyx. It may save catheterization, therefore, if the stretching is done only laterally, i. e., towards the tubera ischii.

When pruritus ani is caused by a local eczema it is well to remember that the latter

RECTUM.

may be seborrheal in origin. In such cases other areas of the disease, as on the chest and, especially, the scalp, should be sought for; they will require attention also, in order to effect a cure.

It is not sufficiently established that the character of the crystals found in the urine indicates the presence or identity of lithiasis in the urinary tract. When cystin crystals are constantly found in the sediment, however, if symptoms of lithiasis are present, the stone is probably made up of cystin.

GENITO- URINARY TRACT.

**Kidney and
Ureter.**

An approximate determination of the origin of a hematuria may be obtained by noting the following points: If pure blood is followed by clear urine, the origin is in the urethra; if the patient first passes urine, then blood, the source of bleeding is probably in the bladder; if urine evenly mixed with blood is voided, the kidney is probably responsible for the hemorrhage; if long, fine clots resembling worms are passed, these, usually, are from the ureter.

A point worth remembering in the diagnosis of nephrolithiasis is that red blood cells

GENITO-URINARY TRACT.

are almost always found in the centrifugalized sediment in the urine even in the interval between attacks of colic.

In a very acid urine red blood cells may be disintegrated and appear under the microscope as an amorphous material. When it is important to determine the presence or absence of blood in the urine it is sometimes necessary, therefore, to resort to a chemical test, e. g., that with guaiac resin.

A radiographic shadow simulating that of a urinary calculus may be produced by an atheromatous plaque, as, for example, in the internal iliac artery, by a phlebolith, or by a calcareous gland.

When skiagraphing for suspected renal calculus the entire urinary tract should be exposed, i. e., the kidney regions, ureters and bladder. Not infrequently a stone supposed to be in the kidney may be lodged in the lower end of the ureter, within reach from the bladder.

By a careful study of an x-ray plate it can usually be determined with fair accuracy

GENITO-URINARY TRACT.

whether a renal calculus is in the pelvis or in the parenchyma, at the upper pole or the lower pole.

In cases of renal colic do not make too positive a pre-operative diagnosis of calculus, no matter how typical the symptoms may be. It has happened very often at the time of operation that no stone is found. Fortunately, these cases are nearly always cured by the exploratory nephrotomy.

Attacks of abdominal pain associated only with intestinal symptoms, may nevertheless be due to a renal or ureteral calculus, even though, in addition, a tender area may be palpated at a point more or less remote from the kidney regions.

The perinephric space is a frequent site of metastatic inflammation after furunculosis or other septic infection.

If possible, always tie each component of a kidney pedicle separately, not en masse.

If a stump ligature, e. g., of the renal pedicle, is slow to come away, the process may

GENITO-URINARY TRACT.

be hastened by fastening it taut to a piece of rubber tubing stretched across the wound.

If pus persists in the urine after the extirpation of a kidney for suppurative disease, it often means that the ureter is involved and will require subsequent extirpation.

When operating upon the ureter for calculus or stricture, avoid undue manipulation; it is important to prevent detachment of the ureter from its bed, if possible.

Pyuria without symptoms is suspicious of an early tuberculosis of the urinary tract.

The examination for tubercle bacilli in the urine by the ordinary method of staining, is not decisive by any means, even if the bladder has been catheterized and differential stains for smegma bacilli have been employed. Numerous examinations with the aid of these procedures must be made, and even then the diagnosis is only a presumptive one. The only sure test is by injecting a large quantity of the sediment into a guinea-pig.

GENITO-URINARY TRACT.

Most cases of sudden, unexpected hemorrhage from the urethra are due to malignant disease, but it is well to remember that there are cases of genito-urinary tuberculosis in which such a hemorrhage is the first symptom.

Bladder.

Hemorrhage from the bladder may yield to irrigations with ice-cold water and with 1-10,000 adrenalin solution, successively.

Never attempt to pack a bladder for hemorrhage without the aid of guy sutures; with them one can make absolutely sure that the gauze goes into the bladder, and not on top of it, pushing the organ away from the space of Retzius.

Post-operative hemorrhage from the base of the bladder that proves inaccessible to ligatures, and uncontrollable by packings, may be checked by the following method: Through several thicknesses of gauze, cut in squares, pass a double strand of heavy silk or of twine fastened on a stout needle. With the patient in Trendelenburg's position and the bladder widely opened, thrust the needle from within directly through the perineum, and bring the gauze firmly against the

GENITO-URINARY TRACT

bleeding surface by pulling upon the threads, which are then to be fastened to an outside dressing.

It is a peculiar fact that many of the cases of tumor of the bladder occur among workers in anilin dyes.

It should be borne in mind that stone in the bladder may be the primary cause in children of enuresis, masturbation or prolapsus recti.

What feels at the other end of the searcher like a stone in the bladder, may be a fold of mucous membrane encrusted with urinary deposit.

In cases of suspected rupture of the bladder, catheterization is not always a sure test. The rent may be so large that the catheter draws away urine that has already flowed into the peritoneal cavity.

To prevent a suprapubic or other drainage tube from becoming displaced is easily accomplished by fitting another tube over it like a collar; this outer tube is split through

GENITO-URINARY TRACT.

half its length and the two portions are spread out over the skin and fastened down with adhesive plaster.

Involuntary urination very often means a distended bladder, and in old men it should at once indicate an examination into the condition of the prostate. Vomiting, too, is often caused by distention of the bladder.

An acutely distended bladder should not be completely emptied in one sitting. Its rapid collapse may produce hemorrhagic cystitis.

Before employing a rubber catheter test its resiliency. If it is brittle or cracked, discard it. Not infrequently a rotten catheter breaks off in the bladder while, of course, a rough catheter or sound may play havoc in the urethra.

Unconscious patients should be catheterized at regular intervals of about eight hours.

Rectal examination sometimes aids in determining the variety of obstruction in prostatic hypertrophy. If the prostate is com-

Prostate.

GENITO-URINARY TRACT.

paratively small, the obstruction is probably due to the middle lobe; if large, to the lateral lobes.

A Mercier catheter is the first kind that ought to be employed in attempting to overcome retention caused by an enlarged prostate. Often it will have to be resorted to in the end; and, therefore, it will save much unsuccessful manipulation to use it at once. Occasionally, a metal catheter will pass when even a Mercier fails.

Never open a prostatic abscess per rectum, no matter how much it bulges; always operate through the perineum.

It is a wise rule to submit all removed hypertrophied prostates to thorough examination by a pathologist. Carcinomatous degeneration may be found in some spot.

Carcinoma of the prostate often does not recur for some time; meanwhile the patient may look surprisingly well. This should not beguile the surgeon into a too hopeful prognosis.

GENITO-URINARY TRACT.

Penis and Urethra.

Force is never helpful in overcoming the resistance of a stricture to instrumental passage; it is bound to do harm. A combination of patience and hot applications, with a strong admixture of gentleness and judgment, will effect the desired result in most cases.

A swelling or redness behind the scrotum, in cases of urethral stricture or developing after urethral instrumentation, usually means urinary extravasation, which requires prompt and active treatment.

One death from urethral sepsis is enough to impress upon one the importance of the teaching that perineal drainage should always be employed after internal urethrotomy three or more inches from the meatus.

When performing external urethrotomy without a guide it is often possible to trace the continuation of the urethra proximal to the opening, by means of a filiform bougie, even when all devices failed to secure the introduction of a filiform before the operation. If a filiform cannot be thus passed through the urethral wound, suprapubic pres-

GENITO-URINARY TRACT.

sure on the bladder may demonstrate the location of the urethral orifice by the escape of a drop of urine or by bulging of the membranous urethra.

Avoid the temptation to employ a constrictor upon the penis when performing circumcision, etc. It may cause sloughing, or actual gangrene.

After circumcision it is important to prevent adhesion of the reflected mucous fold of the prepuce to the corona glandis, by the daily passage of a probe about the corona, and by the use of vaselin.

Absorbent cotton, so commonly used to catch the discharge of gonorrhea, is very inelegant. It sticks to the glans, allows the meatus to glue together, and is difficult to remove without soiling the fingers. The following is the cleanliest and most surgical dressing. In a six inch square of surgical gauze, of about four thicknesses, cut a slit in the middle just large enough to be passed over the glans and to be held behind the corona. Then simply draw the foreskin forward. Indeed, such a dressing will hold

GENITO-URINARY TRACT.

even if the patient has been circumcised, if the slit in the gauze is not too large. With such a simple dressing, there is no retention of the pus, no irritation from contact with the secretions, the organ is readily inspected and the gauze is easily drawn off by a little pull at one of its clean corners.

Non-specific urethral discharges in young boys may be due to foreign bodies introduced while masturbating.

Examine the inguinal regions for hernia in all cases of very tight prepuce causing difficulty in micturition.

If difficulty is experienced in reducing a paraphimosis because of swelling, before dividing the constriction apply a rubber bandage around the parts for a few minutes; this may relieve the swelling to such an extent that the paraphimosis can be easily reduced.

A comforting support for the testicles, when a patient is confined to bed with orchitis, is easily furnished by a well-padded cigar box cover, grooved to fit under the scrotum, and laid across the thighs. Adhesive plaster may be used in the same manner.

**Scrotum
and
Testicle.**

GENITO-URINARY TRACT.

In hydrocele the base of the tumor is below, in spermatocele it is usually above. A milky fluid obtained by aspiration usually speaks for spermatocele.

If a cystic swelling in the scrotum is opaque when examined by the well-known transillumination test, especially if a history of traumatism is elicited, it may still be a hydrocele. Admixture of blood in the hydrocele destroys its translucency.

The early reappearance of fluid after tapping a hydrocele does not necessarily mean that the operation has been a failure. It may be but an inflammatory reaction, subsiding spontaneously or under the application of unguentum iodi.

If a male patient with supposed strangulated hernia complains of pain running down the inner aspect of the thigh it is well to think of torsion of the testicle.

Accumulated experience shows that castration alone will not cure the great majority of cases of tuberculosis testis. In many, if not most, cases the vas deferens,

GENITO-URINARY TRACT.

seminal vesicle or prostate is involved, and it will be necessary to remove one or more of these structures in order to cure. Moreover the other testicle frequently becomes tuberculous. Open-air therapy is helpful.

Syphilitic interstitial orchitis resembles closely in appearance new growth of the testicle. Unless the diagnosis of neoplasm is beyond all doubt, an active course of specific treatment should be tried before removing the testicle.

In excising a varicocele under local anesthesia, tie the upper ligature first; the pain of tying the lower ligature will then be abolished.

After the open operation for varicocele the scrotum may be shortened by simply sewing the wound together transversely instead of longitudinally.

The presence of varicocele, especially if unilateral, should suggest an examination of the abdomen and pelvis for a possible growth pressing on the spermatic veins.

FEMALE GENERATIVE ORGANS.

It is a good rule to always inspect the labia before making a vaginal examination. Many pathological conditions in these parts may otherwise pass unsuspected.

Don't be tempted to exclude gonorrhea because you see no bacterial or other evidence of vaginal or urethral infection. In women the presence of gonorrhea may not make itself known for six weeks or more, and salpingitis may be the first evidence.

When the openings of the Bartholinian glands appear as two sharply defined red spots, an antedating inflammation may be diagnosed with certainty, and in a great majority of instances a latent gonorrhea is present.

Simple incision is not sufficient in the treatment of Bartholinian abscesses. They should be cauterized daily with iodin, and if they recur, excised.

Furunculosis vulvæ, even when it persists in spite of all other treatment, will usually yield to daily scrubbing with green soap and the application of a dressing of sublimate solution.

FEMALE GENERATIVE ORGANS.

When cleansing the vagina and vulva in preparation for an operation, a soft cotton mop should be used for the vestibule; a stiff brush is too apt to bruise or lacerate the urethra and cause dysuria for some days thereafter.

The use of any considerable quantity of iodoformized gauze in the vagina involves the risk of a severe dermatitis of the vulva.

Before performing curettage always make a final bimanual examination of the uterus in narcosis. The finding may determine some other form of treatment. Again, after curettage, before allowing the patient to get out of bed, carefully examine the pelvis for signs of a possible exudate.

As a final cleansing step after curettage of the uterus it is well to introduce, and at once withdraw, a packing of gauze. This brings out with it fragments of tissue not washed out by the irrigation.

Sudden collapse after a curettage for supposed abortion may mean the rupture of an unsuspected ectopic gestation sac.

FEMALE GENERATIVE ORGANS.

Carcinoma of the cervix may remain hidden in the lumen of the cervical canal, which is then eroded and forms an irregular elliptical cavity. While the external os is closed suspicion of the serious condition present will be attracted by the foul or bloody discharge.

No operation for sterility in the female should be performed without first excluding sterility on the husband's part.

In case of hematocolpos and hematometra it is essential to precede all interference by a careful rectal examination in order to determine whether the tubes are distended or not. If hematosalpinx exists a laparotomy and salpingectomy must precede the vaginal operation, otherwise a severe peritonitis may be set up by a reflex discharge of infective secretion from the tubes.

In the early months of pregnancy examinations should be made to determine that there is no retroversion, or to treat it if it exists. A retroverted gravid uterus impacted in the curve of the sacrum always aborts.

Ascites in the presence of a mass in the pelvis usually, but not necessarily, means malignancy.

FEMALE GENERATIVE ORGANS.

Avoid introducing a uterine sound in examinations when pelvic inflammation is suspected. It may set up a parametritis.

Impaction of feces in the sigmoid and rectum, with absorption symptoms, may simulate pelvic peritonitis.

In pulling on the round ligaments in the Alexander operation, use the fingers rather than instruments; a surer hold is given, one can gauge the proper force to employ more readily, and there is less likelihood of the ligaments tearing.

In all cases of lumbago, especially of the chronic variety, examine the sacro-iliac joints for tenderness. Such cases may sometimes be almost instantaneously relieved by applying broad strips of plaster from beyond one superior iliac spine to the other, across the back. The straps must be applied tightly and with the feet close together.

BACK.

A skin-lined sinus opening between the coccyx and the anus, when not very short, usually leads to a dermoid cyst situated close to the coccyx. Frequently loose hairs from the dermoid may be found in the sinus.

BACK.

Congenital paralysis of the lower limbs may arise from an internal sacral or coccygeal spina bifida. In such cases rectal examination reveals the trouble and an operation may afford marked improvement or even a brilliant cure.

EXTREMITIES.

Do not consider too lightly a history of "growing pains" in the extremities in children. These symptoms may be due to a grave osteomyelitis.

Do not be in a hurry to perform primary amputations after severe traumata of the extremities. First, combat the shock and prevent hemorrhage. Keep the wound as clean as possible; and only when the patient has quite recovered from his shock (at the end of a few days or more), perform the amputation.

Pulsation in the course of an artery should not lead to the hasty conclusion that one is dealing with an aneurism. A tumor overlying a large vessel, and also a vascular sarcoma of the bone, may simulate an aneurism very closely.

EXTREMITIES.

Never incise a swelling in the course of a large artery without making sure first that it is not an aneurism.

When clamping a vein in continuity secure the proximal end first; otherwise it will empty and may become lost to view.

In acute (septic) osteomyelitis immediate operation is not too radical; in chronic osteomyelitis patient waiting is often not too conservative—the final expulsion of a sequestrum may be all that is necessary to effect spontaneous cure.

In acute, no less than in chronic osteomyelitis of the long bones, an x-ray picture is of immense service as a guide in the operation. It determines the exact location, extent and even character, of the disease focus, and thus it saves much unnecessary destruction of bone by the surgeon's chisel.

A chronic synovitis of apparently unknown origin and very rebellious to treatment is sometimes due to a small focus of osteomyelitis just beneath the cartilaginous surface.

EXTREMITIES.

When performing amputation, arthrectomy, osteotomy or similar operation it is wiser to leave the constrictor in place until the dressing is partly, or entirely, applied, than to remove it after tying the large vessels, in an effort to secure the small ones. In the former case the snugly applied dressing will safely prevent hemorrhage; in the latter case, there may be an alarming loss of blood from the numerous small vessels in the very time the efforts are made to tie them all.

Do not amputate an extremity for sarcoma without a previous careful examination of the lungs and mediastinum for metastasis. Such symptoms as continued cough, a small hemoptysis or beginning dyspnea, should be regarded as highly suggestive of such a complication.

After major amputations an elastic constrictor should always be left at the head of the bed, so that the nurse can immediately apply it in case of secondary hemorrhage.

Never apply an elastic ligature about the arm without first interposing a towel. This may obviate subsequent paralysis.

EXTREMITIES.

The superficial location of the ulnar nerve must be borne in mind when incising an abscess about the inner aspect of the elbow.

An acute non-purulent tenosynovitis may be satisfactorily treated by immobilization with plaster strips.

Persistent pain in an arm may be due to the presence of a "cervical rib."

Persistent pains in the leg may be due to obliterating endarteritis. This occurs occasionally even in young men and often goes on to the production of gangrene. Both syphilis and excessive smoking are suspected as etiological factors.

Flat-foot is another cause of pains in the leg or thigh.

In cases of pain in the hip of doubtful origin, examination of the kidney regions may discover the cause.

The presence of sciatica demands a careful exploration of the pelvis by rectal or vaginal examination. It should also be re-

EXTREMITIES.

membered that Osler described sciatica as one of the early symptoms of cancer of the breast.

The following are some of the conditions in the presence of which an examination for tabes dorsalis should never be omitted: 1. All primary swellings of the knee or ankle joint without apparent origin. 2. "Sciatica" and "lumbago." 3. A deep ulcer on the base of the great toe. 4. Repeated vomiting at various intervals, with periods of well-being intervening. 5. Abdominal pains without other evident cause.

Pain in the leg after an abdominal operation often means the development of a femoral vein thrombosis. This occurs usually on the left side.

Lymph-edema of the lower extremity associated with a swelling in the groin (fluctuating or not) is significant of carcinoma of the inguinal glands. The primary lesion may be in the rectum, e. g., an epithelioma of the anus that is giving no symptoms.

When removing a lipoma or other growth from the inner surface of the thigh, a little

EXTREMITIES.

care should be exercised in order to avoid cutting the long saphenous vein. Ligature of that vessel (especially in ambulant and in non-aseptic cases) may be followed by a distressing phlebitis.

Inflamed areas and abscesses about the knees of creeping infants should be examined for foreign bodies.

Punctured wounds about the knee should be treated with the greatest solicitude and attention to asepsis, in order to prevent infection of the joint.

In operating for loose bodies within the knee joint, do not be satisfied with removing but one body; a careful examination should be made to determine the presence of more, for they are very frequently multiple.

Do not operate for foreign body in the knee joint without first excluding dislocation of one of the semilunar cartilages.

In amputations below the knee, insist on active and passive motion in the knee joint at an early date. If this is not done contracture ensues, which makes the application of an artificial limb difficult.

EXTREMITIES.

A hematoma may be produced in the calf muscles by direct or indirect violence that the patient may pay little attention to at the time or even fail to recall.

Swelling of the leg, associated with febrile disturbances, may be produced by hematogenous infection of a hematoma of the calf muscles. Such a condition may somewhat simulate osteomyelitis or other serious condition. It may be differentiated, however, by the location of the greatest tenderness and swelling and by a careful inquiry into the history. If no distinct traumatism is recalled the condition of the patient's arteries may nevertheless suggest the possibility of the occurrence of such a hematoma.

Patients with varicose veins should be instructed that in case hemorrhage takes place, the best method of stopping it temporarily is to merely compress the bleeding point with the finger.

Never advise an elastic stocking in cases of varicose veins where recent phlebitis exists. The pressure may detach a part or whole of the thrombus, propelling it into the general circulation.

EXTREMITIES.

Before anesthetizing a patient to operate upon a wound (e. g. of the wrist), in which tendons are severed, attach forceps or ligatures to any tendon ends that are visible. While struggling during primary narcosis the proximal ends of cut tendons are sometimes drawn up, and the above device will obviate slitting up the sheaths to secure them. Squeezing the extremity proximal to the wound will likewise prevent these retractions. **Hand.**

Never divide the annular ligament of the wrist. The hand is much weaker after it is divided than before.

Frequently referred to the surgeon because of the constant pain and marked tenderness, is to be noted a group of cases of what might be termed "occupation wrist pain." They differ from the ordinary case of "writer's cramp," "piano-player's cramp," etc., in that, while these latter frequently have pain in, or about, the wrist, the cases here referred to have no spasm, the pain is constant, and it is not of a neuralgic character. Sometimes it radiates along the thumb (as in mail-openers); sometimes it is localized to the inner border of the lower end of the ulna, which is very sensitive to pressure (as in

EXTREMITIES.

shirt-ironers). The fingers are free. There may be pain in the forearm muscles (flexors).

For a single tenorrhaphy make the incision quite a little to one side of the line of the tendon and perform no more dissection than is necessary. This is to avoid adhesions of the tendon to the skin.

If a tendon has been divided by an incised or lacerated wound and the skin has united over it, it is better to wait a fortnight or more before performing tenorrhaphy. Otherwise organisms introduced with the traumatism may cause suppuration and sloughing of the tendon, not only defeating the operation, but making a later attempt at approximation difficult or impossible.

When exploring for a needle or other foreign body the finger tip is often far more useful than a probe. It must be remembered, too, that strands of fascia often impart to a probe "the feel" of a foreign body. Cutting and picking at these deceptive strands of tissue soon distort the field of operation and destroy important relations. It is extremely desirable to conduct a systematic and

EXTREMITIES.

cleanly dissection when seeking a foreign body.

The best drainage should be afforded for all punctured wounds of the palm; suppurations in this region are very disagreeable and are followed by severe consequences.

The surgeon should not wait for redness before making a diagnosis of palmar abscess. Owing to the density of the fascial structures this sign is often lacking in the early stages.

In dealing with infections of the hand bear in mind that under a simple bleb may lie an extensive phlegmon, threatening, or actually involving, a tendon or bone and urgently needing a generous but wisely placed incision; while on the other hand, a tendon may be thrust from its protecting sheath into the area of destruction by a knife sweep more earnest than judicious. A crater-like opening in a sodden skin, though freely discharging pus, may need enlarging to protect the tissues underlying; while another opening, too long continued by unnecessary packing, may cripple a joint or tendon by undue cicatrization.

In the treatment of hand and finger in-

EXTREMITIES.

fections, it is very important to release from bandaging as much and as many of the fingers as possible, and as soon as possible. The habit of bandaging up immovably all the fingers, in the treatment of a lesion of some of them, saves the surgeon time but, except in short cases, it often cripples the hand by stiffening the fingers.

Occasionally, contractures of the fingers following the treatment of a cellulitis of the hand and forearm may be due, not to the cellulitis itself nor to the incisions made to relieve it, but to fibrosis and shortening of the flexors in the forearm, the result of too tight bandaging or strapping. Such a condition—Volkman's ischemic muscle contracture—must, therefore, be distinguished from the stiff, flexed fingers produced by the cellulitis. Passive motions and massage are helpful in both conditions, but in the former bone shortening (radius and ulna) is necessary to accommodate the contracted muscles.

Remember that chronic ulcers on the hands are found in brass workers, and that a discontinuance of this occupation is necessary to secure healing.

EXTREMITIES.

Indolent sinuses, as of the fingers after deep infections, frequently heal by the daily use of prolonged immersions in hot water.

In dealing with infections or injuries of the fingers amputation should be a last resort. This is especially the case with a thumb, the most important of all the fingers.

In a case of fresh traumatic amputation of a part of the finger, if the amputated part has not been too lacerated or crushed, try to restore the member by cleansing the parts carefully and suturing it to the stump. Once in a while the graft will "take."

Amputation of a finger gangrenous as the result of carbolic acid application should not be performed until the line of demarkation is well established. The necrosis may be superficial and in such an instance the finger may be saved by means of skin graft.

Tenderness in the heel, or pain and tenderness in the sole of the foot is very often, indeed, of gonorrheal origin. It will not be relieved in such cases until treated on that basis. The patient may deny that he ever

Foot.

EXTREMITIES.

had gonorrhea. Examine his urine; shreds tell their own story.

Do not be too hasty in ascribing the cause of pain in the tendo Achilles, or Achilles bursa, to an ill-fitting shoe. First exclude gonorrheal infection.

If the cause of pain in the feet is not otherwise clear, examine them in the dependent position. This may develop the presence of erythromelalgia.

The determination of the presence of a fracture of one of the mid-tarsal bones is extremely difficult, and usually impossible, without x-ray examinations. Yet these examinations have shown the occurrence of such fractures, alone, or associated with injuries to other bones, as the result of injuries by slight or severe direct violence. For this reason, and because fractures of the metatarsals by indirect violence are by no means uncommon, it should be practically a routine to submit the foot to skilful skiagraphy in all cases where either form of violence may have occurred. It will save many patients from weeks of suffering and disa-

EXTREMITIES.

bility. In this region, more than in any other, the x-rays are a means of diagnosis that cannot be dispensed with.

Many, at least, of the sprains of the ankle involve a fracture of the tip of the malleolus, and should be treated by immobilization in plaster-of-Paris.

In old people, as in diabetics, corns, bunions and wounds of the feet demand the most careful attention. They are often the starting points of gangrene.

A very simple method of curing a corn is to excise it.

In ingrowing toenail, evulsion of the nail gives temporary relief, but it does not cure. When the nail grows out again the condition recurs, often in aggravated form.

It is doubtful whether the classical operations for ingrown toe-nail cure permanently in even a fair percentage of cases. Conservative treatment will usually accomplish as much, even in the presence of granulating masses. This treatment includes drawing the

EXTREMITIES.

flesh away from the nail with a strip of adhesive plaster, insertion of a gauze packing under the nail edge and the application of an absorbent antiseptic dressing.

Be very guarded in the prognosis of ulcerations on the sole of the foot in diabetic or tabetic patients, no matter how small or trifling the ulceration may be. They persist for long periods and may even never heal.

FRAC- TURES.

Very often the unskillful treatment of a fracture is worse than no treatment at all. Serious deformities may result from the neglect of small details no less than from the violation of important principles.

The important considerations in the treatment of fractures are, at first, relief of pain and reduction of swelling, and, subsequently, preservation of function of the muscles, the nerves and the neighboring joints. Hence the value of early and frequent massage and passive motion (and in suitable cases, of active motion) and the necessity for avoiding splints that unduly compress the muscles or deprive them of activity.

FRACTURES.

In the treatment of fractures of the forearm no consideration is more important than the avoidance of contractures of the fingers, by the intelligent use of splints and by means of early, active and passive movements.

Permanent contracture of the muscles, notably of the flexor group in the forearm, may develop within a very short time after the application of a splint that exercises undue compression. It is a wise rule to inspect all fracture dressings within twenty-four hours; and when this is not expedient special care should be exercised, when applying the dressing, to avoid compression.

In very many cases it is not necessary to the diagnosis of fracture to elicit crepitus and abnormal mobility—often painful manipulations. In several forms of fracture there are other positive diagnostic evidences. Thus, with Colles' fracture the level of the styloid of the radius will almost always be found to have receded from beyond that of the styloid of the ulna. Moreover, x-ray examinations save much painful manipulation.

After all, the localization of bone tender-

FRACTURES.

ness is not only the most useful sign in determining the site of a fracture, but, even in the absence of other signs, it is often, in itself, diagnostic of the presence of a fracture. As instances may be cited greenstick fracture of the clavicle, and fracture of the metacarpal and metatarsal bones.

In all examinations of children, and in the examination of adults for suspected fractures, leave the painful manipulations for the last.

The x-rays have taught us that mathematical reduction is rarely, and even linear reduction is seldom, accomplished even in cases in which excellent functional results are secured. Radiographs have thus frequently been made the basis of blackmailing damage suits. Nevertheless, the x-rays are, of course, of immense value in the treatment of fractures—not only for reference before and after reduction, but during the reduction itself.

A fracture produced by only slight violence should at once raise the suspicion of a malignant growth. In such a case a uniform dark shadow about the bone as seen

FRACTURES.

in the fluoroscope is to be interpreted as a neoplasm rather than as callus, for recent callus is not opaque to the x-rays.

That a bone appears normal by fluoroscopic examination does not gainsay the presence of a fracture. A fracture of the radius, for example, may occur without displacement of the fragments. An x-ray plate will demonstrate the line of fracture, when the fluoroscope fails to.

Severe localized pain after traumatism, especially in children, may be due to subperiosteal fracture, e. g., near the head of the humerus or the femur. Extreme localized tenderness is the chief sign; abnormal mobility and deformity are absent, and crepitus may not be elicited.

In cases of fracture where an end of the bone lies close beneath the skin do not place a pad or any pressure whatever over this point.

In compound fractures involving loss of continuity do not needlessly remove any piece of bone that has even the smallest attachment.

FRACTURES.

It is surprising how often such pieces heal into the wound and thereby help to save loss of substance.

When applying a plaster dressing to the leg always include the foot if the patient is to be confined to bed; otherwise "drop foot" will develop.

In cases of fracture of a rib, it is necessary to watch the patient carefully for a couple of days to note the onset of a possible lung complication. Localized pneumonitis sometimes occurs.

In severe falls or blows or fractures of the pelvis, catheterize the patient as soon after the injury as possible in order to discover a possible rupture of the bladder.

Fracture of the outer end of the clavicle may follow a fall upon the shoulder. Unless one makes a careful examination such a fracture may escape observation or be mistaken for a dislocation of the outer end of the clavicle, with which condition, indeed, it may be associated.

FRACTURES.

Shortening of the shoulder, as measured from the sternal end of the clavicle to the acromion process, is significant of fracture of the clavicle.

In the aged, pain and disability in the arm after traumatism demand especial care in examination of the shoulder. Fracture of the head of the humerus is often overlooked.

In fractures of the anatomical neck of the humerus, examine carefully for injuries to the brachial plexus.

The radiograph of the elbow of a child shows shadows of numerous epiphyses. One inexperienced with x-ray plates is very apt to mistake one or more of these for fractures. When examining the skiagraph of a child's elbow suspected of fracture or dislocation, it is, therefore, important to have the normal picture in mind, or better yet in hand, for comparison.

Fractures of the head of the radius are probably more common than generally supposed, being overlooked frequently because of the absence of the ordinary signs of fracture.

FRACTURES.

If a small child has been pulled by the arm and thereafter has disability in that member, attention should first be directed to the upper end of the radius. Here one is apt to find a subluxation of the head of the bone ("pulled arm") or an epiphyseal separation.

Marked tenderness over the lower end of the radius, after traumatism, without deformity, is suspicious of fissure of the bone. Mobility or crepitus may be obtainable.

The silver-fork deformity is by no means necessary to the diagnosis of Colles' fracture.

If a patient gives a history of "sprained wrist" that has remained feeble and painful in spite of appropriate treatment for sufficient time, and if the wrist presents thickening and tenderness at its radial aspect, a diagnosis of fracture of the scaphoid should be entertained. Colles' fracture must be excluded, by the relation of the two styloid processes and the location of the deformity. Fractures of the radius and scaphoid may, however, coexist.

FRACTURES.

Fractures of the neck of the femur in old people sometimes cause no other symptoms than disability. The mildness of the trauma and the freedom from much pain should not deceive one.

A padded triangular wooden or cardboard splint—one leg of the triangle bandaged to the thigh, and another to the trunk—makes an excellent ambulatory apparatus in the treatment of fractures of the shaft of the femur in small children. It maintains reduction, leaves the leg free and does not interfere with keeping the child clean.

Cardboard splints can be best molded to an extremity by tearing, instead of cutting them.

In operations for suture of a fractured patella it is very important to sew the torn lateral ligaments of the joint. These aid largely in the support of the joint.

It must be remembered that fractures of the metatarsal bones may be produced by slight injuries. Thus, the base of the fifth metatarsal may be fractured by a twist of the foot while walking or dancing.

LYMPHATIC GLANDS.

An accurate knowledge of the lymphatic drainage of the various regions of the body is absolutely necessary before one can determine the origin of a glandular infection. This is especially important in cancer, when sometimes the glandular involvement offers the first clue to the primary focus.

Do not advise extirpation of large glands in any particular region without making sure that they are not the early manifestations of leukemia or Hodgkin's disease.

If a bubo shows no sign of disappearing under wet dressings, ice bags, etc., and evidences of suppuration are developing, it is better to make a clean dissection and excise the gland without opening it than to incise and drain.

SKIN.

Exposure to the x-rays causes atrophy of the sweat glands; radiotherapy is proving the most satisfactory treatment for hyperhidrosis.

Pure nitric acid, applied on the narrow, blunt tip of a glass rod is successful in the complete destruction of verrucae, but only if it is forced down into their very roots.

SKIN.

Localized, indurated or softening skin infections ("boils") often disappear completely or open painlessly under an application of emplastrum plumbi in which is incorporated 10 per cent. of salicylic acid; or of 10 per cent. to 20 per cent. salicylated soap plaster. After the boil opens the tiny dressing should be changed every two or three hours.

When shaving the hair in the neighborhood of a boil, draw the razor from the base to the apex so as not to drive microorganisms deeper into the tissues.

Do not treat localized subcutaneous red and tender swellings as infections without first making sure that they are not evidences of gout.

Stains of anilin dyes may be removed from the fingers with strong hydrochloric acid, stains of iodine with aqua ammonia, and stains by silver nitrate with potassium iodid solution.

If an incised wound in the soft parts does not heal as readily as it should, examine the urine for sugar.

WOUNDS.

WOUNDS.

In wounds made by coal on the exposed parts of the body, remove all the particles of coal dust; otherwise a disfiguring pigmentation might follow.

A broad clean ulcer on the soft parts often heals per primam if its surface is swabbed with iodin and its edges then brought together with adhesive straps.

An ulcer with indolent flabby granulations may be stimulated to renewed activity by a thorough scraping or by vigorously rubbing it with gauze.

Catgut strands do not always make a good drain for wounds; they tend to swell and occlude.

Fresh wounds about a joint should not be probed to see whether the joint has been penetrated or not. This is an excellent way of infecting it.

The appearance of emphysema in the tissues about an infected wound, accompanied by fever and escape of bubbles of gas from the wound, should be regarded as very omi-

nous, and indicative of gas bacillus infection. Such cases should be treated by extensive incisions.

Blank cartridge wounds must be laid wide open, all dirt and wad carefully removed, and the area swabbed out with tincture of iodine, or with pure carbolic acid followed by alcohol. Tetanus antitoxin should be administered.

If the powder grains are not promptly removed in gunshot wounds of the exposed part, unsightly discolorations result.

The possible development of a duodenal ulcer in cases of extensive burns, must always be borne in mind.

Too prolonged or too rapid and vigorous use of the pump in the Bier apparatus will frequently cause a rupture of the superficial bloodvessels, and in many cases severe sloughing of the superficial parts ensues, the result of the treatment being worse than the primary cause of the trouble. Application of the Bier cup to an abscess for four to five minutes twice a day is more beneficial than a single ten-minute application.

TUMORS.

Soft tumors under the skin, disappearing in the recumbent posture, are usually lymphangiomata.

If a swelling is "fluctuating" do not be too sure that it is not a solid growth. Lymphangiomata fluctuate.

A subcutaneous tumor with a history of puncture or the presence of a minute scar in the overlying skin, usually means that one is dealing with an inclusion or so-called Ranvier cyst.

In hard swellings of doubtful nature marked tenderness is significant of actinomycosis, when acute inflammation may be excluded.

Do not give a good prognosis in cases of melanosarcoma of the fingers or toes, no matter how small the tumor may be, and no matter how high the amputation is performed. In the majority of cases, these patients succumb to metastases.

In the presence of a pulsating tumor, especially of the bone, examine the kidneys. Secondary hypernephromata always pulsate.

TUMORS.

A pulsating tumor of the os ilium (endothelioma, sarcoma) may easily be mistaken for a gluteal aneurism.

The "egg shell crackle" of certain bone tumors is characteristic of multiple myeloma. Examine the urine for albumose.

In cases of bone tumor these organs should never be overlooked in seeking a primary growth—the prostate or mammary glands, according to the sex, and the thyroid.

An amputation for malignant ulceration should not be performed until the possibility of its being merely a broken-down gumma has been satisfactorily excluded.

A metastatic growth in a superficial lymphatic gland or a gland of the skin may sometimes deceptively simulate the appearance of a sebaceous cyst. In a patient suffering with a malignant neoplasm, therefore, the development of a "wen," especially if at an unusual situation, should be regarded with sufficient suspicion to prompt investigation of its interior.

Individuals with bluish sclerotics, and

TUBER- CULOSIS.

TUBERCULOSIS.

with dark lanugo over the upper part of the back, are usually of tuberculous diathesis; and these signs are not inconsequential in making a diagnosis.

Surgical tuberculosis, no less than pulmonary tuberculosis, calls for the most careful general treatment, post-operative and otherwise. Out-of-door life is as important here as for phthisis.

The temptation should not be yielded to to incise a psoas, hip or other "cold" abscess, except in isolated instances and then only under the most rigid asepsis. The production of a mixed infection means chronic sinus, chronic invalidism and, often, amyloid disease.

ANES- THESIA.

In operations upon the head or neck the anesthetist must see to it that no instrument is allowed to lie over the cornea, especially if it is exposed. Ulceration may be caused with ease; it is often healed with difficulty.

During narcosis, when stertorous breathing calls for extension of the jaw, it is well to hold it forward first on one side, then on

the other, alternating at short intervals. Long continued pressure at the angle or angles of the jaw produces much soreness. Often the jaw can be kept forward by catching the lower incisor teeth in front of the upper ones (if they are strong); a single finger on the chin is enough to maintain this position.

In light narcosis the pupils may dilate reflexly from operative manipulations. This, of course, is not to be confused with the sudden extreme dilatation that occurs when the narcosis has been carried too far.

During the conduct of a narcosis, more important than the activity of the conjunctival reflex or the actual size of the pupil in determining the depth of the anesthesia, are the changes in the reactivity of the lid and the alterations in the size of the pupil. They are reliable indices to fluctuations in the depth of the narcosis. Sometimes a patient is quite relaxed and anesthetic although a fair conjunctival reflex is present; and, again, it may occasionally happen that a patient reacts even when that reflex is abolished.

In crying infants it is extremely difficult to determine the presence, and location, of ten-

ANESTHESIA.

der areas. This may be readily accomplished by the administration of chloroform to the extent of primary narcosis. The physical examination then becomes very easy and when a tender spot is handled it will be announced at once by lively reflexes.

During the performance of a hernia operation it is often helpful for the anesthetist to allow the patient to react sufficiently to strain into view a sac that has slipped back into the abdomen.

A convenient way in which the anesthetist may carry, all sterilized and ready for instant use, his hypodermatic solutions, is the following: Shallow, wide-mouthed, half-ounce bottles are sterilized, labeled and filled. Over the mouth of each bottle is then stretched, and hermetically fastened, a cover of sterilized rubber (dam). Before the narcosis is begun the anesthetist disinfects his syringe and sets these bottles in a dish of sublimate solution. This sterilizes the surface of the rubber. When a solution is wanted the needle of the hypodermatic syringe is simply thrust through the rubber and as much as is needed is drawn into the barrel. The puncture hole closes without leak-

age. The covers of the bottles need to be changed only occasionally.

Whenever the arrangement of a patient upon the operating table requires an extremity to occupy a constrained position, that position should be shifted from time to time to avoid pressure paralysis. The anesthetist should never draw the arms alongside the head, nor permit the strap of a leg-holder to press, for more than a few minutes at a time, upon the brachial plexus in the neck.

Nitrous oxid narcosis can, in most cases, be continued "smoothly," with no cyanosis and with fair degree of relaxation, even for an hour. A laparotomy may thus be performed, if ether and chloroform are contraindicated. To secure such a narcosis it is best to use an apparatus that permits exhalation into the gas bag, and which has a valve for the admission of air. The bag should not be distended fully. After brief air and gas administrations, air is turned off and the patient breathes N_2O and his own CO_2 . At short intervals, and whenever there is any cyanosis, a single breath of pure air is allowed.

ANESTHESIA.

Local anesthetics cannot be injected painlessly into tense, inflamed areas unless the injection is begun at a point in the skin well beyond the seat of inflammation.

The admixture of adrenalin to cocain solution counteracts much of the depressant effect of the anesthetic and enhances the local vaso-constriction. When the mixture is used on the surface of a mucous membrane, however, as in excising an ulcer in the mouth, one must be prepared for a marked reactionary bleeding.

INFUSIONS.

For a single intravenous infusion, as to combat the shock of hemorrhage, it is not essential that the solution contain any of the blood salts but the most abundant one—sodium chlorid. For repeated infusions, however, as sometimes used in treating various toxemias, it is better to employ also the other salts, the solution being made of sodium chlorid 0.9, potassium chlorid 0.03, calcium chlorid 0.02, water 100.

Intravenous saline infusions in too large volume are harmful by the production of congestion of the internal viscera. One to one

INFUSIONS.

and a half pints is enough for an adult of average weight.

In performing subcutaneous infusion do not allow too much fluid to accumulate at one area, otherwise necrosis may occur. Shift the needle to various parts not by swinging it from side to side, but by partly withdrawing it and reinserting it to another area.

The pain in the lower part of the back that is so frequently complained of after operation, can be best relieved by placing a small pillow in the hollow of the spine.

POST- OPERATIVE.

In determining the cause of a post-operative fever never fail to look at the throat.

If, after a period of post-operative catheterization, the patient finds herself unable to pass urine spontaneously, apply hot towels to the vulva.

Gastric lavage is the best post-operative anti-emetic.

Vomiting may frequently be controlled by one-drop doses of tincture of iodine in water at half-hourly intervals.

POST-OPERATIVE.

Before putting an unconscious patient to bed, the hot water bags should be removed or sufficiently covered to prevent the occurrence of a burn.

The occurrence of post-operative phlebitis is often encouraged by keeping the patient too long in bed.

Old people should be allowed to sit up or get out of bed as soon after operation as possible in order to avoid post-operative lung complications.

Do not allow patients to lie on the back immediately after an operation involving the vertebræ or the sacrum; a disagreeable necrosis of the skin flaps may rapidly take place.

HEMOR- RHAGE AND SHOCK.

In differentiating shock and concealed hemorrhage progressiveness of the symptoms is very significant of continued bleeding.

Restlessness, increasing pallor, increasing air-hunger, increasing weakness of the pulse, falling temperature (subnormal), and the ephemeral effect of stimulation, all point to

HEMORRHAGE AND SHOCK.

hemorrhage rather than shock. In addition, there is often some local sign or symptom.

In post-operative collapse if, after studying the symptoms, there be any doubt whether the condition be due to shock or to concealed hemorrhage, the wound should be opened and bleeding sought for.

In dealing with secondary hemorrhage from the rectum (whether bleeding vessels are tied or not), it is better to tampon with gauze wrapped about a piece of stout rubber tubing, than with gauze alone.

Bone tenderness, especially of the sternum and tibiæ, is frequently significant of sepsis.

SEPSIS.

In seeking the source of an obscure sepsis, do not overlook an examination of the ischio-rectal region.

Aluminum instruments should not be boiled in soda solution like other instruments. They are to be sterilized by boiling in plain water or by passing them through an alcohol or Bunsen flame.

INSTRUMENTS.

INSTRUMENTS.

Woven catheters may be sterilized by boiling in saturated ammonium sulphate solution. Catheters and bougies may be kept aseptic if they are wrapped in gauze wet with the soap-spirits of the German pharmacopeia.

Warming a laryngeal mirror prevents condensation of the breath upon it for only a short time. The mirror will remain bright, however, throughout a prolonged examination if, instead of warming it, its surface is smeared with an invisible film of soap.

An "invalid table," the shelf of which projects over the patient's body, will be found a great convenience during operations as a receptacle for instruments in immediate use. It saves time and temper, and avoids accumulation of instruments on the patient's body.

When scissors become "catchy" their edges can often be surprisingly smoothed by carrying each blade repeatedly from lock to tip between the firmly pressing thumb and forefinger. Each kind and size of scissors has its own capacity, and should be used

INSTRUMENTS.

only for what it is intended. Ophthalmic instruments are not intended for ordinary dissections, tissue scissors should not be used for cutting bandages, nor bandage scissors for plaster-of-Paris.

Bandage knives cut best when they have a "saw edge," which is easily secured by sharpening them on a window sill or other rough stone.

A scroll-saw, with an assortment of a dozen saws, can be purchased at the hardware store for twenty-five cents; it is ideal for resection of the small bones of the hand and foot, for amputations of the digits, etc.

Well tempered carpenter's chisels and gouges, and a carpenter's wooden mallet answer the purpose admirably for bone work. A useful bone drill can also be selected from the stock of the hardware dealer.

A gardener's pruning knife and a carpenter's miter saw are the best tools for the removal of plaster dressings.

A cheap potato knife, rough sharpened

INSTRUMENTS.

on a stone, is excellent for cutting through starch bandages.

Crochet needles are most useful for lifting buried stitches out of a sinus.

Knitting needles find another purpose as a means of rupturing the membranes when this is needed in obstetrical work.

Sharp and blunt retractors may be fashioned, in an emergency, by bending the tines of a fork and the handle of a spoon, respectively.

A teaspoon is also useful as an elevator of the eye, when resection of the superior maxilla is performed.

An inverted tea-strainer is useful in the dressing after colostomy, to prevent pressure of the gauze upon the gut.

A spoon-shaped potato cutter may be used, in an emergency, as a wound curette.

The multiple surgical uses of the hairpin are also well-known. Of stouter material, if

INSTRUMENTS.

necessary, a small self-retaining retractor can be quickly made from steel wire; it often obviates the need of an assistant when searching the hand or foot for a foreign body.

Similarly, applicators, probes and depressors may be improvised by twisting stout copper wire.

A wedge of hard wood makes a gag quite useful, often, when administering anesthesia.

A discarded thermometer case (or a hard rubber douche point) is a serviceable handle in which to mount, with candle grease or adhesive plaster, a stick of silver nitrate.

A bright and altogether satisfactory light for throat examinations can be had cheaply by covering a 16-candle-power Edison electric bulb with a smooth layer of plaster-of-Paris, about three-eighths of an inch thick, leaving on one side an aperture the size of a silver half-dollar, or larger. The white inner surface of the plaster brilliantly reflects the light. The outer surface may be painted black for appearance's sake.

INSTRUMENTS.

Cheap powder blowers, such as are used for insecticides, may be employed as insufflators in surgical work, and pepper boxes are useful for dusting powders.

Steel spring tape-measures are better than the wires generally sold for the purpose, for conducting to an x-ray tube the current from the coil or static machine; easily kept taut, and quickly adjusted, they are safest for the patient and most convenient for the operator; that they are not insulated is inconsequential—the coverings on the regular wires do not insulate the induced current.

Wooden skewers are serviceable nail-cleaners. Rolling pins and kitchen towel racks are very convenient for adhesive plaster, rubber tissue, etc., especially for hospital dressings.

A probe that has become bent and twisted is readily straightened out by rolling under the foot on an even floor.

SUTURES.

The threading of catgut or kangaroo tendon through a needle-eye not very roomy may be made easy by cutting the suture end

SUTURES.

obliquely and flattening it between the handles of the scissors. Silk must not be cut obliquely, however, for this makes it apt to ravel while it is being threaded.

Silkworm-gut is easily dyed, and incidentally impregnated with an antiseptic, by immersing it for twenty-four hours in one per cent. solution of methyl violet, before the boiling.

When suturing a wound of the scrotum. if the tissue (dartos) is contracted, apply a warm compress for a moment to cause relaxation.

In removing a skin suture, pull up on one side and cut it as close to the skin as possible. This is in order to avoid drawing any of the exposed part of the suture through the wound and thus possibly infecting it.

Everything is to be gained and nothing to be lost by having patients remove enough of their clothing to allow of a completely satisfactory examination in all cases. Instances can be called to mind, by any physician, of erroneous judgments arrived at before exposure of other parts of the body showed con-

EXAMINATIONS.

EXAMINATIONS.

ditions altering one's opinion. Especially is it important to compare the corresponding members of the body on the sound and the affected side, in all doubtful cases.

When a skiagraph shows a condition not recognizable at once as a definite lesion, it is important to make an x-ray picture of the corresponding part of the body on the other side. It may show that the condition is merely a symmetrical peculiarity, and not a pathological one.

Before excluding glycosuria examine both morning and evening specimens of the urine.

When performing an office operation, too great care cannot be taken to sufficiently roll back or remove such articles of clothing as might become soiled. The patient may not say much if he is obliged to draw up a garment wet with blood—but he'll probably think a few things.

Tar-paper is a smooth, fairly waterproof material to tack on the floor when preparing a room for operation.

DRESSINGS. Grocers' paper bags are well-adapted receptacles for soiled dressings.

DRESSINGS.

When a "wet dressing" fails to properly drain a septic wound try a glycerin dressing—gauze wrung out in pure glycerin and covered with waterproof material.

When wet dressings are needed on hairy areas it should not be forgotten that they predispose the hair follicles to infection.

Wet dressings, especially the very useful Burow's solution of aluminum acetate, when applied to the hand or foot, usually cause maceration and whitening of the skin, which is apt to alarm the patient. The addition to the solution of one-fourth its bulk of glycerin or alcohol, will obviate this unsightly maceration.

A bichlorid of mercury dressing should never be applied on an area of skin on which tincture of iodin has been recently painted. An iodid of mercury is formed, which is highly irritating.

Ichthyol, if used in ointment sufficiently strong (25% to 50%), is perhaps the most useful single medicament in aborting early superficial infections.

DRESSINGS.

The addition of a little oil of citronella to an ichthyol ointment robs it of its disagreeable odor.

A 2% ointment of fuchsin in vaselin or zinc oxid frequently yields gratifying results in stimulating the epidermization of indolent ulcers and granulating wounds.

Subiodid of bismuth dusted on an oozing granulating wound promptly stops the bleeding. It is also an excellent stimulant to the growth of epithelium.

An ointment of beta-naphthol, 10; sulphur, 45; lard, 24; and green soap, enough to make 100 parts, is useful in removing gun-powder not too deeply situated in the skin. It must be employed cautiously, however, to avoid a destructive dermatitis.

Gauze is preferable to cotton for padding the axilla or breasts in dressings that are not frequently renewed. Cotton easily becomes matted with sour-smelling secretions and thus sets up dermatitis. The skin over the tendo Achilles and about the heel cannot be too carefully padded, when applying Buck's extension apparatus.

Collodion, commonly used to seal a puncture wound, as after aspiration, will not adhere if the spot is wet or bleeding. To obviate this, pinch up the skin, wipe it dry, apply the collodion and continue the compression a minute or so until the collodion has begun to contract.

When rubber tissue is not at hand to make a "cigarette drain," rubber tubing may be used in its place. Split a piece of tubing of appropriate length, and lay the wick of gauze in the trough thus made, or draw the gauze through the tube with a probe. Fenestrae may be cut as desired.

The painfulness of withdrawing packings that have dried in a wound may be avoided by soaking them with peroxid of hydrogen.

A urethral endoscope will be found a great help as a means of introducing a rubber drainage tube into a narrow, tortuous sinus.

The change of dressings of burns may be made painless, and the growth of epithelium encouraged, by employing next to the wound sterile strips of gutta-percha in the same man-

DRESSINGS.

ner as for skin-grafts. Subiodid of bismuth lightly dusted on the granulating surface stimulates epithelial growth.

Patients will appreciate the use of black bandages for the scalp—where they are comparatively inconspicuous, and for the hands—where they do not soil.

Mastoid and scalp dressing may be reduced in bulk, and the uncomfortable neck turns of the bandage avoided, by the use of starch bandages, which hold neatly and firmly.

Bandages may be fastened in place more neatly and more securely with strips of adhesive plaster, than with safety pins. When bandaging a finger or toe, turns about the hand or foot will be unnecessary if the dressing is fastened down with a narrow strip of plaster run over the top from base to base, and another strip circularly about the dressing at the base of the digit. When using black bandages, employ black adhesive plaster.

INDEX.

	PAGE
Abdomen	25
Anesthesia	90
Appendix	39
Back	61
Bile Tract	30
Bladder	49
Brain	8
Breast	19
 Cranium	 7
 Dressings	 104
 Ear	 8
Examinations	103
Extremities	62
Eye	10
 Face	 12
Foot	73
Fractures	76
 Generative Organs, Female	 58
Genito-Urinary Tract	45
Glands, Lymphatic	84

	PAGE
Hand	69
Head	7
Hemorrhage	96
Hernia	40
Infusions	94
Instruments	97
Intestines	36
Kidney	45
Lymphatic Glands	84
Mouth	13
Neck	14
Nose	11
Penis	53
Pharynx	13
Post-Operative	95
Prostate	51
Rectum	41
Scrotum	55
Sepsis	97
Shock	96
Skin	84
Stomach	32
Sutures	102


INDEX.—*Continued.*

iii

PAGE

Testicle	55
Thorax	21
Tracheotomy	18
Tuberculosis	89
Tumors	88
Urethra	53
Ureter	45
Urinary Tract	45
Wounds	85

COLUMBIA UNIVERSITY LIBRARIES (hsl.stx)
RD 33 B76 1907 C.1
Five hundred surgical suggestions:



2002144743

